

Use of Antipsychotics in Behavioural and Psychological Symptoms of Dementia (BPSD) Discussion Guide

This tool is designed to help providers understand, assess, and manage residents in LTC homes with behavioural and psychological symptoms of dementia (responsive behaviours), with a focus on antipsychotic medications. It was developed as part of Centre for Effective Practice’s Academic Detailing Service for LTC homes. This tool integrates best-practice evidence with clinical experience, and makes reference to relevant existing tools and services wherever possible.

Important principles include:

- Being resident-centred,
- Being mindful of benefits, risks and safety concerns,
- Using an interprofessional team approach and validated tools,
- Prescribing conservatively, and,
- Reassessing regularly for opportunities to deprescribe medications that are no longer needed.

As always, efforts must be made to individualize any treatment decisions for the resident, with consideration given to caregivers, family members, as well as LTC staff.

Identify BPSD Symptom Clusters^{1, 2}

Psychosis



Delusions
Hallucinations
Misidentification
Suspicious

Aggression



Defensive
Resistance to care
Verbal
Physical

Agitation



Dressing/undressing
Pacing
Repetitive actions
Restless/anxious

Depression



Anxious
Guilty
Hopeless
Irritable/screaming
Sad, tearful
Suicidal

Mania



Euphoria
Irritable
Pressured speech

Apathy



Amotivation
Lacking interest
Withdrawn

Overview of BPSD in LTC Management

Treatment for dementia is an ongoing process. Since dementia is a progressive disease, regular follow-ups are necessary to ensure that the resident is receiving the best possible treatment for his or her symptoms. The sections in this tool should each be **considered** at each follow-up (even if some treatments discussed, such as drug therapy, will not be necessary for every resident at every stage of treatment).

Section A: Evaluate BPSD in LTC

Before beginning any sort of treatment (e.g. drug or non-drug therapy), it is important to evaluate the resident's symptoms.

This section discusses:

- Tools for discussing and documenting BPSD
- How to use the P.I.E.C.E.S.TM tools to assess risks to the resident and others
- Clinical evaluations that should take place in order to identify any underlying physiological causes of BPSD



Section B: Initiate Non-Drug Therapy for BPSD in LTC

Non-drug therapy is an important part of managing BPSD, regardless of whether drug therapy is initiated. It is an ongoing process that involves the care team, family, and caregivers.

This section discusses:

- Safety, environmental, and caregiver approach considerations that are core components of non-drug therapy
- Possible solutions to behavioural symptoms, including those identified within the Dementia Observation System (DOS)



Section C: Consider Drug Trial(s)

In some cases, when non-drug therapy approaches alone are not sufficient to manage BPSD, it may be necessary to initiate drug therapy to manage symptoms.

This section discusses:

- Determining the best drug therapy to treat the resident's symptoms
- What symptoms are and are not likely to respond to antipsychotic therapy
- General principles for monitoring, documenting, and following-up on residents receiving medications



Section D: Additional Information on Antipsychotic Therapy

When BPSD are particularly distressing or disturbing, pose an imminent risk of harm to the patient or others, and are likely to respond to antipsychotics (see section C), it is sometimes beneficial to initiate antipsychotic therapy.

This section expands on the information about antipsychotics introduced in Section C, and includes:

- The benefits and harms of antipsychotic therapy
- A table comparing the efficacy of different antipsychotics for treating BPSD, some common side effects, and the cost of treatment
- General guidelines for assessing antipsychotics for possible deprescribing

Section A: Evaluate BPSD in LTC

Remember: Engage the family/caregiver at every step. Discuss any history that may help the care team understand and manage the behaviour (e.g., preferences, activities, routine).

1 Assess & Document

- Document **behaviour or symptom clusters**, including frequency, severity, triggers, and consequences
- Designate specific members of the interprofessional care team who will be responsible for coordinating day-to-day assessment and management
- Standardized clinical assessment tools, such as the Dementia Observation System (DOS)⁴ can be helpful for monitoring and documenting symptoms
- Examples of standardized clinical assessment tools can be found on **Page 7**

2 Identify Risks

- Use the **P.I.E.C.E.S.TM RISKS** mnemonic to assess risks to the resident and others:⁹
 - Roaming:** Is risk greater due to resident roaming?
 - Imminent:** Is significant risk imminent?
 - Suicide:** Does the resident display any suicidal tendencies?
 - Kin:** Is the health or safety of residents/caregivers affected?
 - Self-neglect:** Is resident's self-neglect a risk to themselves or others?

3 Identify BPSD Causes

- Obtain **history** from caregivers, family, and staff¹⁰
- Consider environmental factors and triggers, including possible role of team members
- Consider using **P.I.E.C.E.S.TM** to identify causes (see box on right)

4 Clinical Evaluation¹⁰

The differential diagnosis of the syndrome of behaviour change in dementia is broad. Careful examination of history, physical examination and appropriate investigations may help identify contributing factors. A full, rather than targeted, physical examination is indicated, **within the bounds of patient cooperation.**

History (include family/caregivers):

- Recent changes to environment, routine, sleep pattern, family/social situation
- Medication Review:** Adherence, prescription and OTC medications, anticholinergic load, drugs that may increase agitation (e.g. cholinesterase inhibitors), medication induced hypotension or orthostatic hypotension, medication that may contribute to constipation and urinary retention, drugs and/or alcohol

Physical Examination:

- Be mindful of sources of:**
- Pain (e.g. dental, skin, joint, feet)
 - Hydration (e.g. dehydration)
 - Sensory loss (hearing, vision)
 - CNS change (e.g. new stroke)
 - Infection (e.g. pneumonia, urosepsis)
 - Hypo-perfusion (e.g. new atrial fibrillation, heart failure)
 - Constipation and urinary retention

Laboratory and Imaging (as guided by physical exam/history):

- Blood:** Glucose, calcium, complete blood count (CBC), creatinine, electrolytes, TSH, others as appropriate
- Urine:** Any urinary symptoms? (Note: Caution not to send urine for culture if no urinary symptoms or sudden change in status as "asymptomatic bacteriuria" without lower urinary tract symptoms or symptoms of urosepsis/bacteremia are rarely the cause of increased behavioural symptoms)
- Imaging:** If appropriate (e.g. chest x-ray if suspected pneumonia based on physical exam; CT head if new concerning neurologic findings)

Use P.I.E.C.E.S.TM to Identify Causes⁹

Use the **P.I.E.C.E.S. 3-Question TemplateTM** to ask:

1. What has changed?
2. What are the **RISKS** and possible causes?
3. What is the action?

Consider...

Physical think "the 5 Ds"

Delirium

Disease (cardiovascular, infectious, insomnia, metabolic, nocturia, renal, respiratory, sleep apnea, urinary retention, etc)

Drugs (e.g. acetylcholinesterase inhibitors, anticholinergics, anticonvulsants, anti-Parkinson, benzodiazepines, digoxin, fluoroquinolones, lithium, opioids, systemic corticosteroid)
See Reference List of Drugs with Anticholinergic Effects⁴¹

Discomfort (e.g. pain, constipation, fecal impaction, urinary retention, hunger, thirst)

Disability (e.g. sensory loss)

Intellectual think "the 7 As"

Amnesia (memory)

Aphasia (speech)

Apathy (initiative)

Agnosia (recognition of people or things)

Apraxia (purposeful movement)

Anosognosia (insight/self-awareness)

Altered Perception (sensory information)

Emotional think "the 4 Ds"

Disorder Adjustment (e.g. related to losses)

Disorders of Mood (e.g. depressive symptoms, anxiety)

Delusional (e.g. suspiciousness, psychosis)

Disorders of Personality

Capabilities

Capability too low to meet demands of environment (catastrophic reactions) or not utilized enough (boredom)
Maximize remaining strengths; avoid unnecessary disability

Environment

Consider over-/under-stimulation, relocation, change in routine, noise, lighting, colours, social interactions with caregivers/others

Social

Consider social network, life story, cultural/spiritual heritage

Section B: Initiate Non-Drug Therapy for BPSD in LTC^{11, 12, 13}

Tips for Successful Non-Drug Therapy

- As a general principle, individualize your approach as much as possible. Behavioural triggers and effective ways to treat them will vary from one resident to the next.
- Take advantage of any available system supports, such as Behavioural Supports Ontario (BSO) and Psychogeriatric Resource Consultants (PRCs).¹⁴
- Even if non-drug therapy is successful at managing symptoms (i.e. drug therapy is unnecessary), monitor targeted behaviours for changes and follow-up regularly based on the needs of the resident/caregiver and severity of symptoms.



Safety

- Ensure the resident's safety and other residents' safety by securing the environment
- Make sure you are safe (exit near, chair between you and resident)
- Remove potentially dangerous objects
- Move other individuals away
- Remove ongoing triggers



Environmental Considerations

Eliminate misleading stimuli

- Clutter, TV, radio, noise, people, reflections in mirrors/dark windows, pictures/décor

Reduce environmental stress

- Caffeine, extra people, holiday decorations, public TV

Adjust stimulation

- If over-stimulated, reduce noise, activity, confusion
- If under-stimulated, increase activity/involvement

Enhance function

- Increase lighting, to reduce misinterpretation
- Add signs, cues, or pictures to promote way-finding

Adapt the physical setting according to individual preference

- Secure outdoor areas
- Home-like features
- Smaller, segmented recreational and dining areas
- Spa-like bathing facilities



Caregiver Approach Considerations

Personal approach

- Be calm and compassionate (use/avoid touch as indicated)
- Distract by engaging in individualized activities
- Focus on resident's wishes, interests, concerns
- Approach slowly; look for signs of increase agitation
- Approach resident's private space slowly and ask permission prior to entering

Daily routines

- Keep to the same routine to reduce uncertainty
- Use long-standing history and preferences to guide
- Individualize social and leisure activities to reduce boredom

Communication style

- Most communication is non-verbal, use positive non-verbal cues
- Make eye contact unless perceived as aggressive
- Use short simple words and phrases (residents with dementia have trouble processing multiple words)
- Speak clearly and use a positive tone
- Wait for answers (be patient)

		Behaviour	Possible Solutions
DOS Colours*4		Noisy (Yellow)	<ul style="list-style-type: none"> • Distract, engage • Individualized music, nature sounds, presence therapy (tapes of family)
		Restless (Orange)	<ul style="list-style-type: none"> • Distract, engage • "Rest stations" in pacing path, adapt environment to reduce exit-seeking, physical exercise, outdoor activities
		Exit-seeking (Brown)	<ul style="list-style-type: none"> • Distract, engage • Adapt environment to reduce exit-seeking, physical exercise, outdoor activities • Register the individual with MedicAlert and Alzheimer's Society Safety Home program (contact information will be on bracelet or necklace) • Hide exits with curtains, or paint a black circle on the floor (the individual will think it is a hole and will not exit)
		Verbal aggression (Pink)	<ul style="list-style-type: none"> • Distract, engage • Individualized music, nature sounds, presence therapy (tapes of family)
		Physical aggression (Red)	<ul style="list-style-type: none"> • Distract, keep calm, remain warm and supportive • If possible, give the person some space and try to approach later
Other		Delusion/hallucination	<ul style="list-style-type: none"> • Understand this is their reality and do not confront the false belief • Focus efforts on how the resident feels, not the content; offer distraction, avoid clutter, TV, radio
		Agitated/irritated	<ul style="list-style-type: none"> • Calm, soothe, distract • Individualized music, aromatherapy, pet therapy, physical exercise, outdoor activities
		Resistant to care	<ul style="list-style-type: none"> • Identify source of threat (e.g. pain); change routines and approaches
		Repetitive questions/mannerisms	<ul style="list-style-type: none"> • Reassure, address underlying issue, distract • Put the answer to the same repetitive question on a piece of paper or card and ask the resident to read the card instead
		Hoarding	<ul style="list-style-type: none"> • Remove items gradually, re-organize and clear paths in the case of emergency; be compassionate
		Inappropriate behaviour <small>(e.g. disrobing, masturbation, verbally inappropriate,)</small>	<ul style="list-style-type: none"> • Distract, re-direct • Keep an active and regular schedule to avoid boredom • Try increasing the level of appropriate physical attention • Provide personal space if possible and come back when the resident is calmer • Allow the individual privacy for intimate/personal activities

*DOS = Dementia Observation System (Colours used in table are taken from the DOS system, though you may use different colours in your practice)

Section C: Consider Drug Trial(s)

1 Ensure Drug Trial is Necessary

- Treat underlying causes (e.g. pain, constipation, delirium)
- Ensure that non-drug therapy options have been attempted, and have been unsuccessful

Note: In acute BPSD, if there is a safety risk to patient or others, there may not be time to try non-drug approaches before trying pharmacological management.

2 Select Appropriate Drug Trial

- Select an appropriate drug based on symptoms (see chart at right)
- Identify which behaviour(s) you wish to target (e.g. see symptom clusters on cover page and to right)

- **If you are considering initiating antipsychotic therapy, first ask:**
 - Are symptoms likely to respond to antipsychotics? (see below right)
 - Is there imminent risk of harm to self and/or others?
 - Are symptoms particularly disturbing, distressing or dangerous?
 - Have you weighed the potential benefits and harms? (see page 6)
- See page 6 for a detailed comparison of antipsychotics

- Obtain and document informed consent (see Psychotropic Medication Consent Discussion Tool)¹⁹
- Start with a low dose, and gradually titrate as necessary/tolerated

3 Maintain and Review

- Monitor change in targeted behaviour as well as side effects (see DOS Tool)⁴
- Assess over 1-3 weeks, documenting any benefits and harms realized. If lack of response and/or tolerability, adjust therapy. Increase dose (if not yet maximized) or taper/discontinue¹⁵
- Continue to reassess on an ongoing basis for effectiveness and tolerability
- Consider dose reduction or discontinuation if the drug:
 - Is not effective,
 - Has intolerable side effects, or;
 - Behaviours have been manageable and stable for 3-6+ months¹⁷
- **If considering dose reduction/discontinuation for an antipsychotic, see "Reassessing Antipsychotics for Possible Deprescribing" on page 6**

4 Follow-Up

- Follow-up is important for any drug regimen
- **If antipsychotics used, reassess need at least every 3 months¹⁶**

5 Consider Referral to a Specialist if Drug Trial is Unsuccessful

- If symptoms persist or worsen, consider referral to a specialist

6 Continue Non-Drug Approaches

- Continue using non-drug approaches to prevent further BPSD symptoms








Tips for Drug Trials and Deprescribing

- In all drug trials, unless clinically indicated, start at a low dose and increase or decrease slowly.
- For more tools and resources, visit effectivepractice.org/dementia.
- For more information about antipsychotic deprescribing, including a deprescribing algorithm, visit deprescribing.org.

Selecting an Appropriate Drug Therapy for the Resident's Symptom(s)

Behaviour	Drug Therapy
Psychosis, Aggression, Agitation (severe)	• Atypical antipsychotics (such as risperidone, aripiprazole, olanzapine, quetiapine as discussed in detail on page 6) ^{10,14}
Agitation (severe), unlikely to respond to antipsychotics	• SSRIs such as citalopram or trazodone (however, evidence is lacking for trazodone) ^{15,16,44}
Agitation (severe) in Lewy Body Dementia or Parkinson's	• Possible cholinesterase inhibitors • Very low dose quetiapine ^{15,16}
Anxiety (short term/intermittent)	• A short-acting benzodiazepine such as lorazepam prior to anxiety provoking events such as bathing ¹⁷
Anxiety (chronic)	• Antidepressants (such as SSRIs, SNRIs) • Buspirone ¹⁰
Depression (severe)	• Antidepressants such as SSRIs (e.g., citalopram, sertraline), SNRIs (e.g., venlafaxine, duloxetine), other antidepressants (bupropion, mirtazapine, moclobemide) • Secondary TCAs (nortriptyline or desipramine) may be suitable if coexisting indication like neuropathic pain, etc., but caution regarding anticholinergic load, etc. ^{10,16,18}
Mania	• Addressing any possible drug causes is of primary importance • Evidence for specific recommendations lacking • Mood stabilizers are an option, but take caution regarding tolerability and drug interactions
Apathy	• Limited role for drug therapy but sometimes cholinesterase inhibitors may be helpful • Methylphenidate also sometimes used, but limited by concerns such as stimulant effect on behaviour and risk of diversion ^{15,18}

Symptom Likelihood to Respond to Antipsychotic Therapy

Cluster	Likely	Unlikely
 Psychosis	• Delusions • Hallucinations • Misidentification • Suspicious	
 Aggression	• Defensive • Physical	• Verbal • Resistance to care
 Agitation	• Restless/anxious	• Dressing/undressing • Pacing • Exit seeking ¹⁷ • Repetitive actions ⁴⁵⁻⁴⁷
 Depression	• see below ^{*,**}	• see below ^{*,**}
 Mania	• see below [*]	• Euphoria ⁴⁶⁻⁴⁸ • Irritable ⁴⁶⁻⁴⁸ • Pressured speech
 Apathy^{46,48,49}		• Amotivation • Lack of interest • Withdrawn
 Other		• Hiding or hoarding ⁴⁵ • Wandering without aggression ^{17,45} • Disinhibition (e.g., sexual) ⁴⁵⁻⁴⁷

* The role of antipsychotics in those with dementia and depression is beyond the scope of this evidence review.

** In cases where depression treatment may be indicated, consider psychiatric consultation to determine appropriate pharmacotherapy options.

Section D: Additional Information on Antipsychotic Therapy

Potential Benefits and Harms of Antipsychotic Therapy

Potential benefits tend to be over-appreciated, while harms are underappreciated. Nevertheless, when harmful behaviours are severe and distressing, an antipsychotic trial may be reasonable.

Antipsychotics: Potential Benefits	Antipsychotics: Potential Harms
<p>Limited benefit: modest improvement seldom observed</p> <ul style="list-style-type: none"> • effect size: 0.12-0.2 • NNT variable: ~5-14 <p>(ie. at best, compared to placebo, antipsychotic therapy results in targeted behaviour benefit in 1 out of 5 people treated)^{20,21}</p>	<p>Side effects: sedation, falls, postural hypotension, QT prolongation, confusion, EPS (rigidity, stiffness, akinesia), tardive dyskinesia, diabetes, weight gain^{22,23}</p> <p>Stroke: increased risk</p> <p>Death: possible increase</p> <p>Health Canada Advisory noted a 1.6 fold increase in mortality (mostly related to heart failure, sudden death, pneumonia). Some data suggests that there will be 1 extra stroke or death for every ~100 people treated (NNH=100).^{24, 25, 26}</p>

KEY: EPS: extrapyramidal symptoms (Parkinson's-like); NNT: number needed to treat to see one extra benefit; NNH: number needed to treat to see one extra harm

Comparison of Antipsychotics^{20, 21, 30, 31, 32, 33, 34}

Many effects are dose dependent and direct comparisons are limited. Thus, the following table is intended only as a general guide.

Drug Generic (Brand)	Efficacy or evidence in BPSD therapy	↑BP ³²	Ach	Sedation	EPS	TD ³³	Diabetes	Weight Gain ²⁷	Usual Dose	\$/Month	
Atypicals	Risperidone* (Risperdal) ^{25, 26, 34}	• Indicated for severe dementia of the Alzheimer type (Health Canada) • Evidence for efficacy in agitation, aggression & psychosis	++	++	++	++	+	++	↑↑↑ (0.7lb/month)	0.125mg – 2.0mg/d QHS (or divided BID)	\$10-27
	Olanzapine* (Zyprexa) ^{25, 26, 34}	• Off-label use in BPSD • Evidence for efficacy in agitation & aggression	+	+++	+++	++	+	+++	↑↑↑ (1.0lb/month)	1.25mg – 7.5mg/d	\$17-38
	Aripiprazole* (Abilify) ³⁴	• Off-label use in agitation or aggression ¹⁸ • Evidence for efficacy in agitation & aggression • Not eligible for dementia or BPSD in the elderly ^(ODB criteria, Therapeutic Note) • Not for psychosis ^(same as placebo)	+	+	++	+	+	–	↑	2.0mg – 12.5mg QHS	\$112-260
	Quetiapine (Seroquel) ^{25, 26, 34}	• Off-label use in BPSD • Lacks evidence for efficacy in BPSD agitation, aggression & psychosis • Consider in Lewy Body dementia, Parkinson's (low EPS) • Note: although used, not indicated, and lacking evidence for insomnia	++	+++	+++	+	+	+++	↑↑ (0.4lb/month)	12.5mg – 200mg/d (divided QHS-TID)	\$10-59
Typicals	Haloperidol (Haldol)	• Useful short term in acute BPSD or delirium	+	+	+	+++	+++	++	↑↑	0.25mg – 2.0mg/d	\$14-25
	Loxapine (Loxapac, Xylac) ²	• Consider if other agents have failed and severe, persistent, dangerous behaviour continues • Severe, acute BPSD • Not to be used long-term due to adverse effects	++	++	+++	+++	+++	+	–	5.0mg – 10mg BID	\$18-27

*Aripiprazole, olanzapine and risperidone were superior to placebo as treatment of behavioural symptoms as measured by total scores on BEHAVE-AD³⁶, Brief Psychiatric Rating Scale (BPRS)³⁷, and Neuropsychiatric Inventory (NPI)²⁰

KEY: Terminology

Ach: anticholinergic
 BID: twice daily
 BP: blood pressure
 ODB: Ontario Drug Benefit
 EPS: extrapyramidal symptoms
 lb: pound
 TD: tardive dyskinesia
 TID: three times daily
 QHS: bedtime

Frequency (%) of Adverse Reactions of Antipsychotics at Therapeutic Doses

– : Negligible or absent (<2%)
 + : Infrequent (>2%)
 ++ : Moderate (>10%)
 +++ : Frequent (>30%)
 ↑ : Increase

Tips for Reassessing Antipsychotics for Possible Deprescribing

- Stopping or tapering antipsychotics may decrease “all cause mortality”²⁷
- Deprescribing may not be indicated where symptoms are due to psychosis, or where behaviour is especially dangerous or disruptive
- Evaluate reason for use and any recent changes in targeted behaviour
- Ensure suitable non-pharmacological measures for BPSD are optimized
- Due to the nature of responsive behaviours and the usual course of dementia, antipsychotics can often be successfully tapered and/or discontinued.²⁸ As some may worsen, approach cautiously, and monitor behaviour²⁹
- Taper gradually, often by 25-50% every 2-4+ weeks and look for any resulting behaviour changes. Once on lowest dose, may discontinue in 2-4+ weeks
- Continue to reassess for emergence of responsive behaviours

Supporting Materials

These supporting materials are an inventory for long-term care providers to help identify useful clinical aids and resident/family material. This list includes direct links (where available) to tools or materials that have been directly referenced in the detailing guide and/or have been reviewed by CEP and identified as important material in supporting the uptake/adoption of the detailing guide. It includes a brief description of the tool, with full references available as indicated on **page 8**. The following is a comprehensive but not exhaustive list based on an environmental scan, appraisal by Clinical Leads, and focus groups with long-term care providers. CEP has also compiled a list of additional resources beyond those listed below available at effectivepractice.org/dementia

Assessment and Practice Tools

Antecedent, Behaviour, Consequence (ABC) Chart Form³

Chart form to help providers determine and document the events/stimuli that impact behaviour.

BEHAVE-AD³⁶

Clinical rating scale to measure behavioural and psychological symptoms of dementia based upon information obtained from caregivers/informants.

URL: dementia-assessment.com.au/behavioural

Brief Psychiatric Rating Scale³⁷

Rating scale of 24 symptom constructs used to assess the positive, negative, and affective symptoms of individuals.

Cohen-Mansfield Agitation Inventory (CMAI)⁵

Inventory questionnaire of grouped agitated behaviours to assess the frequency and severity of these behaviours in elderly persons.

Confusion Assessment Method (CAM)⁷

Diagnostic algorithm/questionnaire for identification of delirium through formal cognitive testing.

Cornell Scale for Depression in Dementia³⁸

Scale for assessing signs and symptoms of major depression in people with cognitive impairment.

Kingston Standardized Behavioural Assessment (KSBA)⁶

Behaviour analysis tool designed to indicate the number of behavioural symptoms associated with dementia affecting an individual patient.

URL: kingstonscales.org/behaviour-assessment.html

Neuropsychiatric Inventory – Nursing Home Version (NPI – NH)⁴⁰

Tool to characterize the neuropsychiatric symptoms and psychopathology of patients with Alzheimer's disease and other dementias to measure the impact of antimentia and psychotropic drugs.

URL: npitest.net

Pain Assessment in Advanced Dementia Scale (PAINAD)⁸

Pain assessment tool for individuals with advanced dementia including behaviour observation scores.

Dementia Observation System (DOS)⁴

Behaviour assessment tool which captures the frequency and duration of behaviours of concern over 24 hour periods.

URL: piecescanada.com

PIECES™ Framework⁹

Interdisciplinary approach to understanding and enhancing care for individuals with complex physical/cognitive/mental health need and behaviour changes.

URL: piecescanada.com

Psychotropic Medication Consent Discussion Tool¹⁹

Aid for initiating antipsychotic medications and key discussion items for informed consent from patients or substitute decision makers.

Reference and Support Information

Atypical Antipsychotic Drugs and Dementia – Advisories, Warnings and Recalls for Health Professionals²⁴

Advisory concerning atypical antipsychotic treatment of behavioural disorders in elderly patients, which is associated with an increased risk for all-cause mortality. June 2005

URL: healthycanadians.gc.ca/recall-alert-rappel-avis/hc-sc/2005/14307a-eng.php

Behavioural Supports Ontario (BSO)¹⁴

Integrated network designed to provide services and supports to individuals with behaviours associated with complex mental health, dementia, and other neurological conditions living in long-term care.

URL: brainxchange.ca/Public/Resource-Centre-Topics-A-to-Z/Behavioural-Supports-Ontario.aspx

Meaning and Solutions for Behaviours in Dementia Inventory³⁹

Dementia-related behaviors, including possible causes and solutions for management as a starting point for discussion with caregiver(s).

URL: www.mountsinai.on.ca/care/psych/patient-programs/geriatric-psychiatry

Reference List of Drugs with Anticholinergic Effects⁴¹

Reference list of drugs with low, moderate, and high anticholinergic effects, including side effects and preferred alternatives.

URL: rxfiles.ca/rxfiles

Risperidone - Restriction of the Dementia Indication⁴²

Alert for the restriction of risperidone and related antipsychotic use to patients with severe dementia of the Alzheimer type unresponsive to non-pharmacological approaches and when there is a risk of harm to self or others. February, 2015

Note: *Although recent alert is specific for risperidone, other antipsychotics have similar concerns; however, unlike risperidone, others lack an official indication in BPSD.*

URL: healthycanadians.gc.ca/recall-alert-rappel-avis/hc-sc/2015/43797a-eng.php

Resident Materials

Antipsychotics and Dementia: Discussion Guide

Handout for families, caregivers, and co-residents discussing how antipsychotic medications are used to help people with dementia.

URL: effectivepractice.org/dementia

Treating Disruptive Behaviour in People with Dementia (Patient Material)⁴³

Statements on how to treat disruptive behaviours without antipsychotic drug use.

URL: choosingwisely.org

References

- [1] McShane R. What are the syndromes of behavioral and psychological symptoms of dementia? *Int Psychogeriatr* 2000; 12(Suppl S1):147-153.
- [2] Rabheru K. Practical tips for recognition and management of behavioural and psychological symptoms of dementia. *CGS Journal of CME* 2011; 1(1):17-22.
- [3] Miltenberger RG. Behavior modification: principles and procedure, 4th ed. Belmont, CA: Thomson Wadsworth; 2008.
- [4] Schindel Martin L. The Dementia Observation System: a useful tool for discovering the person behind the illness. *Long-Term Care* 1998; November/December:19-22.
- [5] Cohen-Mansfield J, Marx MS, Rosenthal AS. A Description of agitation in a nursing home. *J Gerontol Med Sci* 1989; 44(3):M77-M84.
- [6] Hopkins RW, Kilik LA. Kingstons Standardized Behavioural Assessment. 2015. Available from: <http://www.kingstonscales.org/behaviour-assessment.html>.
- [7] Inouye S, van Dyck C, Alessi C, Balkin S, Siegal AP, Horwitz R. Clarifying confusion: the confusion assessment method. *Ann Intern Med* 1990; 113(12):941-48.
- [8] Warden V, Hurley AC, Volicer L. Development and psychometric evaluation of the Pain Assessment in Advanced Dementia (PAINAD) scale. *J Am Med Dir Assoc* 2003; 4(1):9-15.
- [9] Hamilton P, Harris D, LeClair K, Collins J. Putting the P.I.E.C.E.S.™ together; a model for collaborative care and changing practice; a learning resource for providers caring for older adults with complex physical and cognitive/mental needs and behavioural Changes, 6th ed. Shop for Learning Publishing Services; 2008. Used with permission P.I.E.C.E.S.™ Canada Consult Group.
- [10] Sorbi S, Hort J, Erkinjuntti T, Fladby T, Gainotti G, Gurvit H et al. EFNS-ENS guidelines on the diagnosis and management of disorders associated with dementia. *Eur J Neurol* 2012; 19(9):1159-79.
- [11] Livingston G, Johnston K, Katona C, et al. Systematic review of psychological approaches to the management of nonpharmacological symptoms of dementia. *Am J Psychiatry* 2005; 162:1996e2021.
- [12] Seitz DP, Brisbin S, Herrmann N, Rappoport MJ, Wilson K, Gill SS et al. Efficacy and feasibility of nonpharmacological interventions for neuropsychiatric symptoms of dementia in long term care: a systematic review. *JAMDA* 2012; 13(6):503-6.e2.
- [13] Vermooij-Dassen M. Meaningful activities for people with dementia. *Aging Ment Health* 2007; 11(4):359-60.
- [14] Behavioural Supports Ontario. Behavioural Supports Ontario (BSO) [Internet]. Available from: <http://brainxchange.ca/Public/Resource-Centre-Topics-A-to-Z/Behavioural-Supports-Ontario.aspx>.
- [15] Herrmann N, Lanctot KL, Hogan DB. Pharmacologic recommendations for the symptomatic treatment of dementia: the Canadian Consensus Conference on the diagnosis and treatment of dementia 2012. *Alzheimer's Res Ther* 2013; 5(Suppl 1):S5.
- [16] Conn D, Gibson M, McCabe D. 2014 CCSMH guideline update - the assessment and treatment of mental health issues in long term care homes: (focus on mood and behaviour symptoms). Toronto: Canadian Coalition for Seniors' Mental Health (CCSMH); 2014.
- [17] Conn D, Gibson M. National guideline for seniors' mental health: the assessment and treatment of mental health issues in long term care homes (focus on mood and behaviour symptoms). Toronto: Canadian Coalition for Seniors' Mental Health (CCSMH); 2006.
- [18] Moore A, Patterson C, Lee L, Vedel I, Bergman H. Fourth Canadian Consensus Conference on the diagnosis and treatment of dementia: recommendations for family physicians. *Can Fam Phys* 2014; 60:433-8.
- [19] Feldman S. Psychotropic medication consent discussion tool. Baycrest Health Sciences: University of Toronto; 2015.
- [20] Maher AR, Maglione M, Bagley S, Suttrop M, Hu JH, Ewing B, et al. Efficacy and comparative effectiveness of atypical antipsychotic medications for off-label uses in adults: a systematic review and meta-analysis. *JAMA* 2011; 306(12):1359-69.
- [21] Maglione M, Ruelaz Maher A, Hu J, Wang Z, Shanman R, Shekelle PG, et al. Off-label use of atypical antipsychotics: an update. *Comparative Effectiveness Review No. 43*. Rockville MD: Agency for Healthcare Research and Quality US; 2011.
- [22] American Geriatrics Society 2015 Beers Criteria Update Expert Panel. American Geriatrics Society 2015 Updated Beers Criteria for Potentially Inappropriate Medication Use in Older Adults. *J Am Geriatr Soc* 2015 Nov; 63(11):2227-46.
- [23] Maust DT, Kim HM, Seyfried LS, Chiang C, Kavanagh J, Schneider LS et al. Antipsychotics, other psychotropics, and the risk of death in patients with dementia: number needed to harm. *JAMA Psych* 2015; 72(5):438-45.
- [24] Health Canada. Atypical antipsychotic drugs and dementia – advisories, warnings and recalls for health professionals [Internet]. 2005. Available from: <http://www.healthycanadians.gc.ca/recall-alert-rappel-avis/hc-sc/2005/14307a-eng.php>.
- [25] Schneider LS, Tariot PN, Dagerman KS, Davis SM, Hsiao JK, Ismail MS, et al. Effectiveness of atypical antipsychotic drugs in patients with Alzheimer's disease. *NEJM* 2006; 355:1525-38.
- [26] Schneider LS, Dagerman KS, Insel P. Risk of death with atypical antipsychotic drug treatment for dementia: meta-analysis of randomized placebo-controlled trials. *JAMA* 2005; 294(15):1934-43.
- [27] Ballard C, Hanney ML, Theodoulou M, Douglas S, McShane R, Kossakowski K, et al. The dementia antipsychotic withdrawal trial (DART-AD): long-term follow-up of a randomized placebo-controlled trial. *Lancet Neurol* 2009; 8:151-57.
- [28] Declercq T, Petrovic M, Azermi M, Vander Stichele R, De Sutter AI, van Driel ML et al. Withdrawal versus continuation of chronic antipsychotic drugs for behavioural and psychological symptoms in older people with dementia. *Cochrane Database Syst Rev* 2013; 28:3.
- [29] Pan YJ, Wu CS, Gau SS, Chan HY, Banerjee S. Antipsychotic discontinuation in patients with dementia: a systematic review and meta-analysis of published randomized controlled studies. *Dement Geriatr Cogn Disord* 2014; 37(3-4):125-40.
- [30] Muench J, Hamer A. Adverse effects of antipsychotic medications. *Am Fam Phys* 2010; 81(5):617-22.
- [31] Taylor D, Paton C, Kapur S. Prescribing guidelines in psychiatry, 11th ed. Wiley-Blackwell; 2012.
- [32] Bezchlibnyk-Butler KZ, Jeffries JJ, Procyshyn RM, Virani AS, ed. Clinical handbook of psychotropic drugs, 17th ed. Hogrefe & Huber Publishers; 2007.
- [33] bpacnz. Prescribing atypical antipsychotics in general practice. *BPJ* 2011; 40:14-23.
- [34] Pharmacists Letter, Prescribers Letter. Comparison of Atypical Antipsychotics PL Detail-Document #281006. Stockton, CA: Therapeutic Research Center; October, 2012.
- [35] University of Iowa College of Public Health. Antipsychotic medicines for people with dementia [Internet]. Iowa: University of Iowa; 2013. Available from: https://www.healthcare.uiowa.edu/IGEC/IAAdapt/view/Antipsychotic_consumer.pdf.
- [36] Adapted from Reisberg B et al. Behavioral symptoms in Alzheimer's disease: phenomenology and treatment. *J Clin Psych* 1987; 45(5 Suppl):9-15.
- [37] Overall, JE, Gorham DR. The brief psychiatric rating scale (BPRS): recent developments in ascertainment and scaling. *Psychopharmacol Bulletin* 1988; 24:9-99.
- [38] Alexopoulos GS, Abrams RC, Young RC, Shamoian CA. Cornell scale for depression in dementia. *Biol Psychiatry* 1998; 23:271-84.
- [39] Danielli E. Meaning and Solutions for Behaviours in Dementia Inventory. [Internet] Toronto: Mount Sinai Hospital; 2013. Available from: <http://www.mountsinai.on.ca/care/psych/patient-programs/geriatric-psychiatry/prc-dementia-resources-for-primary-care/dementia-toolkit-for-primary-care/responsive-behaviours-in-dementia>.
- [40] Cummings JL. Neuropsychiatric inventory – nursing home version (NPI – NH) [Internet]. 1994. Available from: <http://www.npistest.net>.
- [41] Bareham J. Anticholinergics: reference list of drugs with anticholinergic effects [Internet]. RxFiles; 2014. Available from: <http://www.rxfiles.ca/rxfiles/uploads/documents/Psych-BPSD-Newsletter.pdf>.
- [42] Health Canada. Risperidone - restriction of the dementia indication [Internet]. 2015. Available from: <http://healthycanadians.gc.ca/recall-alert-rappel-avis/hc-sc/2015/43797a-eng.php>.
- [43] Choosing Wisely Canada. Canadian Geriatrics Society: Treating disruptive behaviour in people with dementia: antipsychotic drugs are usually not the best choice [Internet]. Choosing Wisely Canada; 2014. Available from: <http://www.choosingwiselycanada.org/materials/treating-disruptive-behaviour-in-people-with-dementia-antipsychotic-drugs-are-usually-not-the-best-choice/>.
- [44] Porsteinsson AP, Drye LT, Pollock BG, Devanand MD, Frangakis C, Ismail Z, et al. Effect of citalopram on agitation in Alzheimer disease: the CitAD randomized clinical trial. *JAMA* 2014; 311(7): 682-691.
- [45] Brodaty H, Ames D, Snowden J, Woodward M, Kirwan J, Clarnette R, et al. A randomized placebo-controlled trial of risperidone for the treatment of aggression, agitation, and psychosis of dementia. *J Clin Psych* 2003; 64(2): 134-43.
- [46] Paleacu D, Barak Y, Mirecky I, Mazeh D. Quetiapine treatment for behavioural and psychological symptoms of dementia in Alzheimer's disease patients: a 6-week, double-blind, placebo-controlled study. *Int J Geriatr Psychiatry* 2008; 23: 393-400.
- [47] Streim JE, Porsteinsson AP, Breder CD, Swanink R, Marcus R, McQuade R, et al. A randomized, double-blind, placebo-controlled study of aripiprazole for the treatment of psychosis in nursing home patients with Alzheimer disease. *Am J Geriatr Psychiatry* 2009; 16(7): 537-50.
- [48] De Deyn PP, Rabheru K, Rasmussen A, Bocksberger JP, Dautzenberg P, Eriksson S, et al. A randomized trial of risperidone, placebo, and haloperidol for behavioral symptoms of dementia. *Neurology* 1999; 53: 946-55.
- [49] Tariot PN, Schneider L, Katz IR, Mintzer JE, Street J, Copenhaver M, et al. Quetiapine treatment of psychosis associated with dementia: a double-blind, randomized, placebo-controlled clinical trial. *Am J Geriatr Psychiatry* 2006; 14(9): 767-76.

This Tool was developed by Centre for Effective Practice (CEP) with input and review from our Academic Detailing Service Team, including our clinical leads Drs. Andrea Moser and Sid Feldman. In addition, target end-users and other relevant stakeholders provided input during the development process. This Tool was funded by the Province of Ontario as part of CEP's Academic Detailing Service. For more information on the Academic Detailing Service, our Team and/or the development process please visit: effectivepractice.org/academicdetailing.

This Tool was developed for licensed health care professionals in Long-Term Care as a guide only and does not constitute medical or other professional advice. Physicians, Nurse Practitioners, Pharmacists and other health care professionals are required to exercise their own clinical judgment in using this Tool. Neither the Centre for Effective Practice ("CEP"), nor any of their respective agents, appointees, directors, officers, employees, contractors, members or volunteers: (i) are providing medical, diagnostic or treatment services through this Tool; (ii) to the extent permitted by applicable law, accept any responsibility for the use or misuse of this Tool by any individual including, but not limited to, Long-Term Care providers or entity, including for any loss, damage or injury (including death) arising from or in connection with the use of this Tool, in whole or in part; or (iii) give or make any representation, warranty or endorsement of any external sources referenced in this Tool (whether specifically named or not) that are owned or operated by third parties, including any information or advice contained therein.



Use of Antipsychotics in Behavioural and Psychological Symptoms of Dementia (BPSD) Discussion Guide: Long-Term Care (LTC) Edition is a product of the Centre for Effective Practice. Permission to use, copy and distribute this material for all non-commercial and research purposes is granted, provided the above disclaimer, this paragraph and the following paragraphs, and appropriate citations appear in all copies, modifications, and distributions. Use of the Use of Antipsychotics in Behavioural and Psychological Symptoms of Dementia (BPSD) Discussion Guide: Long-Term Care (LTC) Edition for commercial purposes or any modifications of the tool are subject to charge and use must be negotiated with the Centre for Effective Practice (info@effectivepractice.org).

For statistical and bibliographic purposes, please notify the Centre for Effective Practice (info@effectivepractice.org) of any use or reprinting of the tool. Please use the below citation when referencing the tool:

Reprinted with Permission from Centre for Effective Practice. (April 2016). Use of Antipsychotics in Behavioural and Psychological Symptoms of Dementia (BPSD) Discussion Guide: Long-Term Care (LTC 2nd Edition).

Developed by:

